



RESEARCH ARTICLE SUMMARY

MICROBIOME

Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*

M. A. Hanson*, L. Grollmus, B. Lemaitre*

INTRODUCTION: Antimicrobial peptides (AMPs) are host-encoded immune effectors first characterized for their role in fighting infection. AMPs are also important in determining the composition of the host microbiome in both plants and animals. Although many studies have shown rapid evolution of AMPs, little is known about the selective pressures driving that evolution.

RATIONALE: The host microbiome should exert a substantial selective pressure on host immune molecules because the host must maintain a delicate balance with its microbial associates. Variation in a single AMP can upset this balance, as suggested by recent investigations across diverse taxa. In *Drosophila*, previous studies have shown the AMP family *Diptericin* (*Dpt*) evolves rapidly, including a major effect of the amino acid polymorphism S69R of *DptA* on host defense against the opportunistic pathogen *Providencia rettgeri*, and *Providencia* spp. are commonly found in fly

microbiome communities. Beneficial bacteria of the host microbiome also grow out of control in flies lacking multiple AMP gene families, particularly the gut mutualist *Acetobacter*. *Drosophila* species encode two *Diptericin* genes, *DptA* and *DptB*, which are the product of an ancestral duplication stemming from a *DptB*-like gene. To test the idea that the host immune repertoire might be specifically evolved for controlling common microbiome bacteria, we screened recently made *Drosophila* AMP mutants for defense against infection by *Acetobacter* spp. to determine whether any of the AMP genes could explain how flies keep this mutualistic microbe under control.

RESULTS: We found that a single AMP gene, *DptB*, explains the host ability to resist infection by multiple *Acetobacter* species. This interaction is highly specific: We confirmed that *DptA* does not contribute to defense against *Acetobacter*, whereas *DptB* does not contribute to defense

against *P. rettgeri*. We therefore determined the evolutionary history of the *Diptericin* locus, and performed a systematic review of microbiome literature of *Drosophila* and other Diptera. We realized that there have been at least two events of convergent evolution toward *DptB*-like genes in flies feeding on fruit, an ecology associated with high levels of *Acetobacter*. These observations suggest that *DptB* evolved to control *Acetobacter* in the fruit-feeding *Drosophila* ancestor. Moreover, flies that secondarily adopted a mushroom-feeding ecology have repeatedly lost their *DptB* genes, alongside an absence of *Acetobacter* in mushroom-breeding sites. A similar pattern of evolution is also seen in flies that have developed a plant-parasitic ecology, which have lost both *DptA* and *DptB* genes and have an ecology lacking both *Providencia* and *Acetobacter*. To investigate whether these AMP-microbe specificities are shared throughout *Drosophila*, we infected species from across the phylogeny with a diverse complement of *DptA*- and *DptB*-like genes and alleles. We included species with a diversity of *DptA*-like genes, and both *Drosophila melanogaster* and mushroom-feeding flies with or without *DptB*. Host resistance to infection by *P. rettgeri* and *Acetobacter* was readily predicted using just *DptA* or *DptB* presence and polymorphism status, even across fly species separated by about 50 million years of evolution.

CONCLUSION: Our study shows how two microbe-specific defences evolved due to an ancestral duplication producing two *Diptericin* genes. We describe a one-sided evolutionary dynamic wherein the host has adapted its immune repertoire to environmental microbes rather than coevolution of host and microbe. This finding helps to explain the evolutionary logic behind the bursts of rapid evolution common in AMP gene families across taxa. Our results also reveal why certain AMPs can have such disproportionate roles in defense against specific microbes: They were evolutionarily selected for that purpose. This realization suggests that the genome can encode “vestigial” immune effectors, AMPs evolved for defense against microbes that are no longer relevant to the host’s modern ecology. Thus, derivation and loss of microbe-specific effectors offers the immune system a highly effective mechanism for tailoring host defenses for control of ecologically relevant microbes. ■



Fruit fly experiments demonstrate that the host immune system is uniquely adapted to common environmental microbes. Evolutionary selection can tailor host antimicrobial peptides (chains) to control specific microbiome bacteria. As a defense system common across plants and animals, variations in the repertoire of antimicrobial peptides are likely important as key risk factors for preventing infection by common ecological microbes. [Credit: Diego Galagovsky]

The list of author affiliations is available in the full article online.

*Corresponding author. Email: m.hanson@exeter.ac.uk (M.A.H.); bruno.lemaitre@epfl.ch (B. L.)

Cite this article as M. A. Hanson et al., *Science* 381, eadg5725 (2023). DOI: 10.1126/science.adg5725

S READ THE FULL ARTICLE AT
<https://doi.org/10.1126/science.adg5725>

RESEARCH ARTICLE

MICROBIOME

Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*M. A. Hanson^{1,2*}, L. Grollmus¹, B. Lemaitre^{1*}

Antimicrobial peptides are host-encoded immune effectors that combat pathogens and shape the microbiome in plants and animals. However, little is known about how the host antimicrobial peptide repertoire is adapted to its microbiome. Here, we characterized the function and evolution of the *Diptericin* antimicrobial peptide family of Diptera. Using mutations affecting the two *Diptericins* (*Dpt*) of *Drosophila melanogaster*, we reveal the specific role of *DptA* for the pathogen *Providencia rettgeri* and *DptB* for the gut mutualist *Acetobacter*. The presence of *DptA*- or *DptB*-like genes across Diptera correlates with the presence of *Providencia* and *Acetobacter* in their environment. Moreover, *DptA*- and *DptB*-like sequences predict host resistance against infection by these bacteria across the genus *Drosophila*. Our study explains the evolutionary logic behind the bursts of rapid evolution of an antimicrobial peptide family and reveals how the host immune repertoire adapts to changing microbial environments.

Animals live in the presence of a complex network of microorganisms known as the microbiome. The relationship between host and microbe can vary from mutualist to pathogen, which is often context dependent (1). To ensure presence of beneficial microbes and prevent infection by pathogens, animals produce many innate immune effectors as a frontline defense. Chief among these effectors are antimicrobial peptides (AMPs), small, cationic, host defense peptides that combat invading microbes in plants and animals (2–5). Although many studies have shown important roles for AMPs in regulating the microbiome [reviewed in Bosch and Zasloff (6)], presently, we cannot determine why animals have the particular repertoire of AMPs that their genome encodes.

Innate immunity has been characterized extensively in *Drosophila* fruit flies (7, 8). Antimicrobial peptide responses are particularly well characterized in this insect (2, 9, 10). In *Drosophila*, AMP genes are transcriptionally regulated by the Toll and Imd nuclear factor- κ B (NF- κ B) signaling pathways (8). Recent work has shown that individual effectors can play prominent roles in the defense against specific pathogens (11–19). Consistent with this, population genetics studies have highlighted genetic variants in AMPs correlated with susceptibility against specific pathogens. A landmark study in *Drosophila* found that a serine-arginine

polymorphism at residue 69 in one of the two fruit fly *Diptericins*, “S69R” of *DptA* (Fig. 1A), is associated with increased susceptibility to *Providencia rettgeri* bacterial infection (20). A loss-of-function study later showed that flies lacking both *Diptericin* genes (“*Dpt*^{SK1},” flies lacking *DptA* and *DptB*) are as susceptible to *P. rettgeri* infection as Imd pathway mutants, whereas flies collectively lacking five other AMP families nevertheless resist infection in a manner similar to the wild type (21). Like these investigations in *Drosophila*, a G49E polymorphism in the AMP Calprotectin of Persian domestic cats is associated with susceptibility to severe ringworm fungal skin disease (22). Similar AMP variation is common across animals (23–26). However, although *P. rettgeri* is an opportunistic pathogen of wild flies and ringworm is common in certain cat breeds, whether these AMPs are evolving to selection imposed by these microbes is unclear. Given recent studies on AMP roles beyond infection (27–31), other fitness trade-offs could also explain AMP evolution.

It is now clear that antimicrobial peptides shape the microbiome (6), but defining if or how the host immune repertoire itself is shaped by the microbiome has been challenging. Here, we characterized the function and evolution of the *Diptericin* gene family of flies, revealing that these AMPs were selected to control ecologically relevant microbes.

Results

Diptericin B is specifically required for defense against *Acetobacter* bacteria

Acetobacter bacteria are mutualists of *Drosophila* that supplement host nutrition and are common in wild flies (32–35). We previously showed that a strain of *Acetobacter* grows out of control in

the gut of *Relish* mutant flies (*Rel*^{E20}) lacking Imd pathway activity and in flies carrying deletions removing 14 AMP genes (Δ AMP14) (36). Here, we identified this *Acetobacter* species as *A. sicerae* strain BELCH (fig. S5). Gnotobiotic association with *A. sicerae* did not cause mortality, even in Δ AMP14 flies (fig. S6A). However, pricking flies with a needle contaminated with *A. sicerae* killed Δ AMP14 flies (12, 36), also causing an abdominal bloating phenotype that preceded mortality (shown later). This route of bacterial infection is similar to what flies experience when their cuticle is pierced by natural enemies [e.g., nematodes, wasps, and mites (37–39)]. Because Δ AMP14 flies are killed by *A. sicerae* systemic infection, one or more AMPs are likely required to control opportunistic infections by this microbe. We therefore used flies carrying overlapping sets of AMP mutations (21), including a *Diptericin* mutant panel affecting each of the two *Diptericins* (Fig. 1B), to narrow down which AMP(s) protects the fly against *A. sicerae* infection.

Ultimately, deleting just *DptB* fully recapitulates the susceptibility of Δ AMP14 flies. *Dpt*^{SK1}, *DptB*^{KO}, and *DptB*^{A3} flies suffered 100% mortality after infection, with survival curves mirroring Δ AMP14 and *Rel*^{E20} flies; these *DptB*-deficient flies also presented similar levels of abdominal bloating (Fig. 2, A and B). Furthermore, ubiquitous RNA interference (RNAi) silencing of *DptB* caused both mortality and bloating after *A. sicerae* pricking (fig. S6, B and C). Conversely, *DptA*^{S69R}, *DptA*^{A622}, and even Δ AMP8 flies collectively lacking five other AMP gene families [*Drosocin*, *Attacin*, *Defensin*, *Metchnikowin*, and *Drosomycin* (21)] resisted infection in a manner comparable to wild type. Finally, *DptB* mutants display increased *A. sicerae* loads, preempting mortality (Fig. 2C), suggesting a direct role for *DptB* in suppressing *A. sicerae* growth.

After revealing the critical importance of *DptB* in defense against *A. sicerae*, we investigated whether *DptB* has a broader role in the control of other *Acetobacter* species. To this end, we infected flies with a panel of *Acetobacter* species including *A. aceti*, *A. indonesiensis*, *A. orientalis*, *A. tropicalis*, and *A. pomorum*. Although these *Acetobacter* species displayed different levels of virulence, *DptB* specifically promoted survival and/or prevented bloating against all virulent *Acetobacter* species (figs. S7 and S8).

Collectively, these results indicate that *DptB* is an AMP of specific importance in defense against multiple *Acetobacter* species, revealing another example of high specificity between an innate immune effector and a microbe relevant to host ecology. Because *Acetobacter* are common in fermenting fruits (40, 41), the major ecological niche of *Drosophila*, *DptB* might be especially important for flies to colonize this niche.

¹Global Health Institute, School of Life Science, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland. ²Disease Ecology and Evolution, Biosciences, University of Exeter, Penryn, United Kingdom.

*Corresponding author. Email: m.hanson@exeter.ac.uk (M.A.H.); bruno.lemaitre@epfl.ch (B.L.)




Fig. 1. Dipterics of *D. melanogaster*. (A) Alignment of *D. melanogaster* mature DptA and DptB peptides, which are ~52% identical. The DptA^{S69R} site is noted (Q in DptB, and see fig. S1 for protein folding predictions). (B) The two *Diptericin* genes are located in tandem on chromosome 2R:55F with only 1130 base pairs (bp) between them. DptA^{Δ822} encodes a premature stop (W40*). Strain DptB^{Δ3} encodes a 37-bp deletion overlapping the DptB intron-exon boundary, causing loss of function (fig. S2). The Dpt^{SK1} deficiency removes 2137 bp, deleting the coding region of both genes. DptB also encodes a secreted propeptide (PP), similar to *Drosophila* Attacins (figs. S1, S3, and S4). SP, signal peptide.




Fig. 2. DptB is specifically required for defense against *A. sicerae*. (A) Flies lacking DptB bloat after *A. sicerae* systemic infection. Each data point reflects the average from one replicate experiment (~20 males). (B) Sum survival curves showing that DptB is critical for defense against *A. sicerae*. (C) *A. sicerae* bacterial load increases before mortality. Each data point reflects the average of five pooled flies. *n*^{exp}, number of experiments.

Diptericin A is specifically required to defend against *P. rettgeri*

The Gram-negative bacterium *P. rettgeri* was isolated from the hemolymph of wild-caught flies (20), suggesting that it is an opportunistic pathogen in *Drosophila*. Previous studies showed that *Diptericins* play a major role in surviving *P. rettgeri* infection (20, 21), including a marked correlation between the DptA S69R polymorphism and resistance against this bacterium: Flies encoding arginine were more susceptible than flies encoding serine at this site (20). However, it is unknown if DptB contributes to defense against *P. rettgeri*.

We therefore infected our panel of *Diptericin* mutants by pricking with *P. rettgeri* (Fig. 3A). We confirmed the DptA^{S69R} allele reduces survival after *P. rettgeri* infection, here with a controlled genetic background ($P < 2 \times 10^{-16}$). DptA^{Δ822} flies also paralleled mortality of Dpt^{SK1} flies lacking both *Diptericin* genes ($P = 0.383$). Initially, we found that DptB^{KO} flies showed higher susceptibility to *P. rettgeri*

($P = 9.44 \times 10^{-11}$), correlated with higher bacterial load (fig. S9A). However, our isogenic DptB^{KO} flies had only ~57% induction of the DptA gene compared with our isogenic DptA^{S69} wild type at 7 hours after infection (fig. S9B). By contrast, we observed that DptB^{Δ3} flies carry the DptA^{S69} allele, have wild-type DptA expression (fig. S2), and actually survive infection by *P. rettgeri* even better than DptA^{S69} ($P = 5.03 \times 10^{-4}$; Fig. 3A). Moreover, silencing DptB by RNAi did not significantly affect survival against *P. rettgeri* ($P > 0.05$; Fig. 3B). We therefore conclude that DptB itself does not have a major effect on resistance to *P. rettgeri*, although a cis-genetic background effect found in DptB^{KO} flies causes lesser induction of DptA and, accordingly, higher susceptibility.

Our *Diptericin* mutant panel shows that DptA plays a major role in defense against *P. rettgeri* but not *A. sicerae*. Conversely, DptB plays a major role against *A. sicerae* but not *P. rettgeri*. Thus, these two *Diptericin* genes are highly specific effectors explaining most

of the Imd-mediated defense of *D. melanogaster* against systemic infection by either bacterium.


The Dipterics family shows multiple bursts of rapid evolution across Diptera

Given the high specificity of *D. melanogaster* *Diptericins* for different ecologically relevant microbes, we next investigated whether host ecology might explain *Diptericin* evolution. First, we reviewed the evolutionary history of *Diptericins* across Diptera using newly available genomic resources (Fig. 4).

Diptericins are found across brachyceran fly species, indicating an ancient origin of this antibacterial peptide (>150 million years ago) (42, 43). The extant *Drosophila* DptB-like gene was originally derived in the Drosophilidae ancestor through rapid evolution [figs. S11 and S12; first shown in (43, 44)]. Later, a duplication of DptB gave rise to the DptA locus in the Drosophilinae ancestor ~50 million years ago [date per (45)], which began as a DptB-like

Fig. 3. *DptA* is specifically required for defense against *P. rettgeri*.

(A) Sum survival curves of *Diptericin* mutants after infection with *P. rettgeri*. (B) Silencing *DptB* by RNAi (*Act>DptB-IR*) does not significantly affect fly survival compared with *Act>OR* controls (RNAi validation is shown in fig. S10).



gene but then evolved rapidly after the duplication [shown in (44); also see figs. S11 and S12 and table S1]. Given these repeated bursts of evolution and only ~52% similarity between *DptA* and *DptB* (Fig. 1A), distinct antibacterial activities are not necessarily surprising. In reviewing *Diptericin* evolution, we further realized the *DptA*^{S69} residue of *D. melanogaster* is also present in the subgenus *Drosophila* through convergent evolution: Different codons are used by the subgenus *Sophophora* (e.g., AGC) and subgenus *Drosophila* (e.g., TCA) to produce *DptA*^{S69} residues (table S1), providing further evidence that adaptive evolution selects for serine at this site [complementing (20) and (44)]. Moreover, across species, there is a high level of variation at this site: In addition to the S69R polymorphism, this site can also encode either glutamine (Q) or asparagine (N) in *DptA* of other *Drosophila* species. Q/N is also seen at the aligned residue of *DptB* across *Drosophila* species (Q56N in *DptB*). These four residues (S, R, Q, and N) are derived compared with the ancestral aspartic acid residue (D) found in most other dipterans (table S1).

This analysis suggests that the extant *DptB*-like gene first evolved in the drosophilid ancestor, whereas *DptA* emerged from a duplication of a *DptB*-like gene, followed by rapid diversification. The *DptA*^{S69} residue was also derived at least twice, and this site is highly polymorphic across genes and species. These repeated bursts of evolution suggest that fly *Diptericins* evolved responding to selection in the drosophilid ancestor.

Diptericin evolution correlates with microbe presence in host ecology

The diversity of *Drosophila* ecologies, along with many wild-caught fly microbiome studies, places us in a unique position to pair each host's microbial ecology with patterns in the evolution of their *Diptericins*, which have microbe-specific importance.

We performed a systematic review of the Diptera microbiome literature (table S2). *Acetobacter* bacteria are regularly found across species feeding on rotting fruits in microbiome

studies (32, 34, 46, 47). However, *Acetobacter* appear to be absent from rotting mushrooms (48), and are largely absent in wild-caught mushroom-feeding flies themselves (48, 49). Further, *Providencia* bacteria related to *P. rettgeri* are common in species feeding on both rotting fruits and mushrooms [(34) and table S2]. We observed that three drosophilid species with mushroom feeding ecology, *D. testacea*, *D. guttifera*, and *Leucophenga varia*, have independently lost their *DptB* genes (Fig. 4) (43). Thus, three independent *DptB* loss events have occurred in flies with a mushroom-feeding ecology specifically lacking in *Acetobacter*.

There is another *Drosophila* sublineage with an ecology that lacks *Acetobacter*: *Scaptomyza* (Fig. 4, green branch). *Scaptomyza pallida* feeds on decaying leaf matter and mushrooms, whereas *Scaptomyza flava* and *Scaptomyza graminum* feed on living plant tissue as leaf-mining parasites (50). The *S. flava* microbiome shows little prevalence of either *Acetobacter* or *Providencia* (51). We investigated whether these *Scaptomyza* species had pseudogenized either of their copies of *DptA* (two genes, *DptA1* and *DptA2*) or *DptB* (one gene). We found independent premature stop codons in *DptA1* in the leaf-mining species *S. flava* (Q43*) and *S. graminum* (G85*), but not in the mushroom-feeding *S. pallida* (fig. S13). We also analyzed the promoter regions of these *DptA* genes for the presence of Relish NF-κB transcription factor binding sites ["Rel-κB" sites from (52); fig. S13A], confirming that the *S. pallida* *DptA1* promoter retains Rel-κB sites and likely immune induction. Thus, *Scaptomyza* *DptA1* genes show pseudogenization specifically in the leaf-mining species that lack *Providencia* in their present-day ecology. However, *DptA1* appears functional in *S. pallida*, a mushroom-feeding species likely exposed to *Providencia* through its ecology. *Scaptomyza* *DptA2* genes show variable presence of Rel-κB sites, but no obvious loss-of-function mutations in coding sequence, and *DptA2* remains expressed in *S. flava* (fig. S13B). Screening the *DptB* genes of *Scaptomyza*, we found no obvious loss-of-function mutations in

coding sequences. However, all three *Scaptomyza* species lack Rel-κB sites in their *DptB* promoter regions (fig. S13A). Whether due to plant feeding or mushroom feeding, none of these *Scaptomyza* have an ecology associated with *Acetobacter*. Using RNA-sequencing data from the *S. flava* midgut (53), we confirmed a lack of expression of both the pseudogene *DptA1* and *DptB* compared with the abundant expression of *DptA2* (fig. S13B). We conclude that *Scaptomyza* species have independently pseudogenized *DptA* and *DptB* genes correlated with presence or absence of *Providencia* or *Acetobacter* in their ecology.

Finally, convergent evolution toward *DptB*-like sequence has occurred in another lineage of "fruit flies": Tephritidae (43, 44) (see figs. S11 and S12 for protein alignment and paraphyly of tephritid *Diptericins* clustering with drosophilid *DptB*). This family of Diptera is distantly related to Drosophilidae (last common ancestor ~111 million years ago). Like *Drosophila*, many tephritid lineages (e.g., Trypetinae and Dacinae) feed on fruits, but like *Scaptomyza*, one lineage, Tephritinae, parasitizes live plants (Fig. 4, purple branches). In light of the present study, it would seem that the tephritid species that feed on *Acetobacter*-associated fruit (40, 54, 55) have convergently evolved a *DptB*-like gene, including a parallel Q/N trans-species polymorphism at the critical Diptericin residue (table S1). Like *Scaptomyza*, plant-parasitizing tephritids lack both *Acetobacter* and *Providencia* in their microbiomes (43) and have lost their *Diptericin* genes (Fig. 4) (43). Thus, *DptB*-like genes evolved in both Tephritidae and Drosophilidae species associated with a fruit-feeding ecology in which *Acetobacter* is a dominant member of the microbiome. The fact that *DptB*-like genes are not found in species unless their ancestor had a fruit-feeding ecology suggests two things: (i) that the *Acetobacter*-rich fruit-feeding niche was colonized before the derivation of *DptB*-like sequence and (ii) that selection imposed by *Acetobacter* resulted in the ancestors of both Tephritidae and Drosophilidae evolving *DptB*-like genes to help control this microbe.




Fig. 4. Dipterin evolution correlates with host ecology and presence of *Acetobacter* or *Providencia*. Dipterin presence was screened in diverse Diptera. The residue aligned to the *DptA*^{S69R} or *DptB*^{Q56N} polymorphism is shown. The *DptB*-like sequence evolved first in the ancestor of Drosophilidae, and the serine-coding allele in *DptA* evolved at least twice (fig. S11 and table S1). The relatedness of the codons used to encode the S/R/Q/N polymorphism enables their diversification in the subgenus *Sophophora* (summary in top left). Fruit-feeding tephritids convergently evolved a *DptB*-like gene (figs. S11 and S12) including a parallel Q/N polymorphism, and *P. variegata* encodes an independent *DptB* duplication, in which the two daughter genes encode either version of the Q/N polymorphism. Within Drosophilidae (bottom part), three species with mushroom-feeding ecology have lost their *DptB* genes: *L. varia*, *D. testacea*, and *D. guttifera*. In both Drosophilinae (*Scaptomyza*) and Tephritidae (Tephritinae), divergence to plant feeding is also correlated with loss of Dipterin genes (fig. S13). Systematic review of microbiome studies (table S2) suggests that the absence of *Providencia* and *Acetobacter* in the host ecology is correlated with *DptA* and *DptB* loss, respectively. Red [x] indicates that the gene loss was confirmed. Copy number variation is noted in table S1. Phylogenetic cladogram was drawn from consensus of multiple studies (45, 65–67).

Our phylogenetic and ecological survey reveals multiple parallels among the host immune effector repertoire, ecology, and the associated microbiome. This suggests that these dipteran species have derived *DptA*- or *DptB*-like genes as their evolutionary solution to control important bacteria found in their microbiome. By contrast, specific Dipterin genes become superfluous when their hosts shift to ecologies lacking Dipterin-relevant microbes, leading to gene loss.

Variation in *DptA* or *DptB* predicts host resistance across species separated by 50 million years of evolution

Our study indicates that among the suite of immune genes involved in *Drosophila* host defense, the AMPs *DptA* and *DptB* are critically important against two environmentally relevant bacteria: the opportunistic pathogen *P. rettgeri* and the gut mutualist *Acetobacter*. Moreover, our phylogeny-microbiome analysis reveals substantial correlations in terms of gene

emergence, retention, and loss. If *DptA* and *DptB* really evolved to control *P. rettgeri* and *Acetobacter*, then the outcomes of *P. rettgeri* and *A. sicerae* infection across species should be readily predicted using just variation in these two Dipterins. We therefore chose 12 *Drosophila* species with variation in the polymorphic site in *DptA* and presence or absence of *DptB*, and infected them with *P. rettgeri* or *A. sicerae*. Experiments in *D. melanogaster* suggest that *DptA*^{S69R} affects defense against *P. rettgeri*, but how *DptA*^{S69Q} or *DptA*^{S69N} affects defense against this bacterium has never been tested. Similarly, the effect of *DptB*^{Q56N} on defense is also untested, so we have no a priori expectations for how these polymorphisms affect peptide activity. To analyze these experiments, we used a linear mixed-model approach (see the materials and methods), including *D. melanogaster* flies from our Dipterin mutant panel as experimental controls. This helped to calibrate our model for the expected effect size for variants of *DptA* or *DptB* within a single species or across species. We also conducted these experiments at 21°C to avoid heat stress to some species, which reduced *D. melanogaster* mortality compared with 25°C (fig. S14).

Summaries of fly species mortality are shown in Fig. 5. As found in *D. melanogaster*, resistance to *P. rettgeri* was associated with a *DptA*^{S69R} allele across species. Indeed, *DptA*^{S69R} found in either *D. melanogaster* or *D. willistoni* correlates with increased susceptibility to *P. rettgeri* ($t = -9.59$, $P < 2 \times 10^{-16}$). *Drosophila yakuba* with *DptA*^{S69N} was also more susceptible than its close relatives, suggesting that asparagine (N) is an immune-poor allele against *P. rettgeri* ($t = -7.26$, $P = 4 \times 10^{-13}$). Further, *DptA*^{S69Q} flies (*D. suzukii* and *D. immigrans*) had similar survival after *P. rettgeri* infection compared with *DptA*^{S69R} flies ($t = +0.07$, $P = 0.35$), suggesting that glutamine (Q) is a competent defense allele against *P. rettgeri* when coded by *DptA* (Fig. 5A). Overall, ~74% of variation in susceptibility can be attributed to variation in *DptA* alone as a fixed effect (marginal $R^2 = 0.743$).

For infections with *A. sicerae*, the absence of *DptB* in the mushroom-feeding species *D. testacea* and *D. guttifera* was correlated with increased susceptibility compared with their close relatives ($t = -10.83$, $P < 2 \times 10^{-16}$). Mushroom-feeding flies displayed increased susceptibility to *A. sicerae* infection that was independent of *DptB* status ($t = -3.77$, $P = 2 \times 10^{-4}$). However, even within this susceptible lineage, *DptB* loss still increased mortality to a similar extent as *DptB* deletion in *D. melanogaster*, indicating that the contribution of *DptB* to defense against *A. sicerae* is independent of host genetic background (Fig. 5B). Overall, ~87% of variation in susceptibility to *A. sicerae* can be explained by just *DptB*




Fig. 5. Dipteriscins predict pathogen-specific survival across *Drosophila*. Host phylogeny, ecology, and *Diptericin* complement are shown. Clean injury is shown in fig. S15. **(A)** Susceptibility to *P. rettgeri* infection varies across species, with survival largely explained by the *DptA* allele, particularly within the subgenus *Sophophora* (blue-shaded species). **(B)** Susceptibility to infection by *A. sicerae* is predicted by presence or absence of *DptB*, although mushroom-feeding flies also had a higher susceptibility to *A. sicerae* infection that was independent of *DptB* loss. Each data point represents one replicate experiment using 20 male flies.

absence and host ecology as fixed effects (marginal $R^2 = 0.868$).

These survival data establish that the specific resistance conferred by *Diptericins* observed in *D. melanogaster* applies across *Drosophila* species separated by ~50 million years of evolution. We conclude that the host immune repertoire adapts to the presence of ecologically relevant microbes through the evolution of specialized AMPs as weapons to combat specific microbes.

Discussion

Susceptibility to infection often correlates with host phylogeny (56, 57), although host ecology greatly influences microbiome community structure (34, 58). Early studies of immune evolution suggested that AMPs were mostly generalist peptides with redundant function, suggesting that AMP variation was not caused by adaptive evolution (2, 3). Instead, studies on immune adaptation have found whole pathway-level effects or have identified factors specific to a given species [e.g., host-symbiont coevolution (59–61)]. As a result, despite a rich literature on immunity-microbiome interactions, the evolutionary logic explaining why the host genome encodes its particular immune effector repertoire has been difficult to approach experimentally.

Here, we identified how ecological microbes promote the rapid evolution of effectors of the immune repertoire, tailoring them to be highly microbe specific. The two *D. melanogaster*

Diptericin genes also provide a textbook example of how gene duplication can promote immune novelty, equipping the host with extra copies of immune tools that can be adapted to specific pathogen pressures. The *Drosophila* *Diptericin* mechanism of action has been elusive because of technical difficulties in peptide purification (2, 10). However studies using *Phormia terranova* highlight many directions for future research [(42, 62, 63) and see discussion in the supplementary materials]. Future studies combining both fly and microbiome genetics should be fruitful in learning how host and microbiome factors determine specificity. One goal of infection biology is to try to identify risk factors for susceptibility present in individuals and populations. Our study suggests that characterizing the function of single effectors, interpreted through an evolution-microbe-ecology framework, can help to explain how and why variation after infection occurs within and between species.

The fly *Diptericin* repertoire reflects the presence of relevant microbes in that species' ecology. Conversely, loss or pseudogenization of *Diptericins* is observed when the microbes they target are no longer present in their environment. In a sense, this means that some AMPs seen in the genomes of these animals are vestigial: Immune genes evolved to fight microbes that the extant host rarely encounters (e.g., *DptB* in *D. phalerata*). Indeed, flies that lack *DptB* genes are likely disadvantaged on *Acetobacter*-rich food resources, where the

possibility of *Acetobacter* systemic infection poses a constant threat. Thus, loss of this AMP makes recolonization of *Acetobacter*-rich rotting fruits a risky proposition, entrenching the host in its derived ecological niche.


Although other mechanisms of defense surely contribute to resistance, *Diptericins* have evolved recurrently as the fly genome's solution to control specific bacteria. Given our findings, we propose a model of AMP-microbiome evolution that includes gene duplication, sequence convergence, and gene loss, informed by the host ecology and the associated microbiome (Fig. 6). In doing so, we thus explain one part of why various species have the particular repertoire of AMPs that they do. This ecology-focused model of AMP-microbiome evolution provides a framework for understanding how host immune systems rapidly adapt to the suite of microbes associated with a new ecological niche. These findings are likely of broad relevance to immune evolution in other animals.

Methods summary

Full materials and methods are found in the supplementary materials. In brief, *D. melanogaster* fly stocks included both natural mutations and a transgenic insertion disrupting *DptB*, which were isogenized into the DrosDel isogenic background, as indicated in Fig. 1 with the prefix “iso.” Nonisogenic *DSPR A3* flies (*DptB^{Δ3}*) were from (64). Survival experiments were performed and analyzed as described

Fig. 6. AMP evolution explained using *Diptericin* and a microbial ecology framework.

Outgroup ecologies per Fig. 4. (1) The drosophilid ancestor fed on fruit and was consequently exposed to *Acetobacter* bacteria. The *DptB*-like sequence evolved rapidly ($dN > dS$) to control this new microbe. (2) A duplication of *DptB* gave rise to the *DptA* locus. (3) The addition of a second *Diptericin* gene permitted evolutionary tinkering to control another relevant microbe, *P. rettgeri*, including convergent evolution of the critical S69 residue. In the sublineage including *D. melanogaster*, codon volatility enables any of S, R, Q, or N residues. The sublineage including *D. guttifera* and *S. flava* evolved its S residue using a different codon, evolutionarily fixing this residue (table S1). (4) In *D. melanogaster*, host ecology remains associated with both *Acetobacter* and *Providencia*, which continually select for maintenance of both genes. (5) In mushroom-feeding *D. guttifera*, *Providencia* remains a threat, but mushroom ecology lacks *Acetobacter*. Consequently, selection is relaxed on *DptB*, leading to pseudogenization. (6) In leaf-mining species such as *S. flava*, *Acetobacter* and *Providencia* are absent from the microbiome, and thus selection is relaxed on both *Diptericin* genes. This AMP-evolution-ecology framework makes sense of why AMPs have microbe specificity and helps to explain how shifts in microbial ecology can promote rapid evolution for AMP-microbe specificity or loss of “vestigial AMPs” that are relevant primarily against microbes that the host no longer encounters.



previously (21), with the temperature and optical density at 600 nm (OD_{600}) of the bacteria (“OD”) indicated within figures. Twenty male flies were used per experiment unless otherwise indicated, and at least three replicate experiments were performed for all data shown in main figures, with raw data available in the supplement. In fly bloating, bacterial load, and gene expression graphs, error bars indicate SD. The cladogram and annotations in Fig. 4 were generated by literature review (table S2), with gene search and annotation methods per (43).

The script for Fig. 5 is available in the supplement. Briefly, we used a linear mixed-model (“lme4” and “performance” packages in R) with species relatedness and experiment block included as random factors and host ecology and variation in *DptA* or *DptB* loci including copy number or alleles at key residues (*D. melanogaster* *DptA* N52 or S69 alleles) as fixed factors. When loss of function was present, the allele was called as “deleted.” We explored our model both by Akaike information criterion model selection and by iterative linear mixed-model testing in which nonsignificant fixed factors (e.g., *DptB* allele in explaining sur-

vival after *P. rettgeri* infection) and their interactions were relegated to being random factors in the final model. These two approaches provided similar results, and we used values from linear mixed models in the main text.

REFERENCES AND NOTES

- J. H. Daskin, R. A. Alford, Context-dependent symbioses and their potential roles in wildlife diseases. *Proc. Biol. Sci.* **279**, 1457–1465 (2012). doi: [10.1098/rspb.2011.2276](https://doi.org/10.1098/rspb.2011.2276); pmid: [22237907](https://pubmed.ncbi.nlm.nih.gov/22237907/)
- M. A. Hanson, B. Lemaître, New insights on *Drosophila* antimicrobial peptide function in host defense and beyond. *Curr. Opin. Immunol.* **62**, 22–30 (2020). doi: [10.1016/j.coi.2019.11.008](https://doi.org/10.1016/j.coi.2019.11.008); pmid: [31835066](https://pubmed.ncbi.nlm.nih.gov/31835066/)
- B. P. Lazzaro, M. Zasloff, J. Rolff, Antimicrobial peptides: Application informed by evolution. *Science* **368**, eaau5480 (2020). doi: [10.1126/science.aau5480](https://doi.org/10.1126/science.aau5480); pmid: [32355003](https://pubmed.ncbi.nlm.nih.gov/32355003/)
- N. Mookherjee, M. A. Anderson, H. P. Haagsman, D. J. Davidson, Antimicrobial host defence peptides: Functions and clinical potential. *Nat. Rev. Drug Discov.* **19**, 311–332 (2020). doi: [10.1038/s41573-019-0058-8](https://doi.org/10.1038/s41573-019-0058-8); pmid: [32107480](https://pubmed.ncbi.nlm.nih.gov/32107480/)
- S. Hacquard, S. Spaepen, R. Garrido-Oter, P. Schulze-Lefert, Interplay Between Innate Immunity and the Plant Microbiota. *Annu. Rev. Phytopathol.* **55**, 565–589 (2017). doi: [10.1146/annurev-phyto-080516-035623](https://doi.org/10.1146/annurev-phyto-080516-035623); pmid: [28645232](https://pubmed.ncbi.nlm.nih.gov/28645232/)
- T. C. G. Bosch, M. Zasloff, Antimicrobial peptides—Or how our ancestors learned to control the microbiome. *mBio* **12**, e0184721 (2021). doi: [10.1128/mBio.01847-21](https://doi.org/10.1128/mBio.01847-21); pmid: [34579574](https://pubmed.ncbi.nlm.nih.gov/34579574/)
- T. B. Sackton et al., Dynamic evolution of the innate immune system in *Drosophila*. *Nat. Genet.* **39**, 1461–1468 (2007). doi: [10.1038/ng.2007.60](https://doi.org/10.1038/ng.2007.60); pmid: [17987029](https://pubmed.ncbi.nlm.nih.gov/17987029/)
- B. Lemaître, J. Hoffmann, The host defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* **25**, 697–743 (2007). doi: [10.1146/annurev.immunol.25.022106.141615](https://doi.org/10.1146/annurev.immunol.25.022106.141615); pmid: [17201680](https://pubmed.ncbi.nlm.nih.gov/17201680/)
- S. J. H. Lin, L. B. Cohen, S. A. Wasserman, Effector specificity and function in *Drosophila* innate immunity: Getting AMPed and dropping Boms. *PLoS Pathog.* **16**, e1008480 (2020). doi: [10.1371/journal.ppat.1008480](https://doi.org/10.1371/journal.ppat.1008480); pmid: [32463841](https://pubmed.ncbi.nlm.nih.gov/32463841/)
- J.-L. Imler, P. Bulet, Antimicrobial peptides in *Drosophila*: Structures, activities and gene regulation. *Chem. Immunol. Allergy* **86**, 1–21 (2005). doi: [10.1159/000086648](https://doi.org/10.1159/000086648); pmid: [15976485](https://pubmed.ncbi.nlm.nih.gov/15976485/)
- L. B. Cohen, S. A. Lindsay, Y. Xu, S. J. H. Lin, S. A. Wasserman, The Daisho peptides mediate *Drosophila* defense against a subset of filamentous fungi. *Front. Immunol.* **11**, 9 (2020). doi: [10.3389/fimmu.2020.00009](https://doi.org/10.3389/fimmu.2020.00009); pmid: [32038657](https://pubmed.ncbi.nlm.nih.gov/32038657/)
- M. A. Hanson, S. Kondo, B. Lemaître, *Drosophila* immunity: The *Drosocin* gene encodes two host defence peptides with pathogen-specific roles. *Proc. Biol. Sci.* **289**, 20220773 (2022). doi: [10.1098/rspb.2022.0773](https://doi.org/10.1098/rspb.2022.0773); pmid: [35730150](https://pubmed.ncbi.nlm.nih.gov/35730150/)
- M. A. Hanson et al., The *Drosophila* Baramicin polypeptide gene protects against fungal infection. *PLoS Pathog.* **17**, e1009846 (2021). doi: [10.1371/journal.ppat.1009846](https://doi.org/10.1371/journal.ppat.1009846); pmid: [34432851](https://pubmed.ncbi.nlm.nih.gov/34432851/)
- M. Shaka, A. Arias-Rojas, A. Hrdina, D. Frahm, I. Iatsenko, Lipopolysaccharide-mediated resistance to host antimicrobial peptides and hemocyte-derived reactive-oxygen species are the major *Providencia* alcalifaciens virulence factors in *Drosophila melanogaster*. *PLoS Pathog.* **18**, e1010825 (2022). doi: [10.1371/journal.ppat.1010825](https://doi.org/10.1371/journal.ppat.1010825); pmid: [36084158](https://pubmed.ncbi.nlm.nih.gov/36084158/)
- C. Zanchi, P. R. Johnston, J. Rolff, Evolution of defence cocktails: Antimicrobial peptide combinations reduce mortality and persistent infection. *Mol. Ecol.* **26**, 5334–5343 (2017). doi: [10.1111/mec.14267](https://doi.org/10.1111/mec.14267); pmid: [28762573](https://pubmed.ncbi.nlm.nih.gov/28762573/)
- R. Xu et al., The Toll pathway mediates *Drosophila* resilience to *Aspergillus* mycotoxins through specific Bomanins. *EMBO Rep.* **24**, e56036 (2022). pmid: [36322050](https://pubmed.ncbi.nlm.nih.gov/36322050/)
- J. I. Perlmuter, J. R. Chapman, M. C. Wilkinson, I. Nevarez-Saenz, R. L. Unckless, A single amino acid polymorphism in natural

- Metchnikowin alleles of *Drosophila* results in systemic immunity and life history tradeoffs. *bioRxiv* 2023.01.16.524277 [Preprint] (2023). <https://doi.org/10.1101/2023.01.16.524277>.
18. A. W. Clemmons, S. A. Lindsay, S. A. Wasserman, An effector peptide family required for *Drosophila* toll-mediated immunity. *PLoS Pathog.* **11**, e1004876 (2015). doi: [10.1371/journal.ppat.1004876](https://doi.org/10.1371/journal.ppat.1004876); pmid: [25915418](https://pubmed.ncbi.nlm.nih.gov/25915418/)
 19. J. Huang *et al.*, A Toll pathway effector protects *Drosophila* specifically from distinct toxins secreted by a fungus or a bacterium. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2205140120 (2023). doi: [10.1073/pnas.2205140120](https://doi.org/10.1073/pnas.2205140120); pmid: [36917667](https://pubmed.ncbi.nlm.nih.gov/36917667/)
 20. R. L. Unckless, V. M. Howick, B. P. Lazzaro, Convergent balancing selection on an antimicrobial peptide in *Drosophila*. *Curr. Biol.* **26**, 257–262 (2016). doi: [10.1016/j.cub.2015.11.063](https://doi.org/10.1016/j.cub.2015.11.063); pmid: [26776733](https://pubmed.ncbi.nlm.nih.gov/26776733/)
 21. M. A. Hanson *et al.*, Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *eLife* **8**, e44341 (2019). doi: [10.7554/eLife.44341](https://doi.org/10.7554/eLife.44341); pmid: [30803481](https://pubmed.ncbi.nlm.nih.gov/30803481/)
 22. A. N. Myers *et al.*, An ancient haplotype containing antimicrobial peptide gene variants is associated with severe fungal skin disease in Persian cats. *PLoS Genet.* **18**, e1010062 (2022). doi: [10.1371/journal.pgen.1010062](https://doi.org/10.1371/journal.pgen.1010062); pmid: [35157719](https://pubmed.ncbi.nlm.nih.gov/35157719/)
 23. R. L. Unckless, B. P. Lazzaro, The potential for adaptive maintenance of diversity in insect antimicrobial peptides. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371**, 20150291 (2016). doi: [10.1098/rstb.2015.0291](https://doi.org/10.1098/rstb.2015.0291); pmid: [27160594](https://pubmed.ncbi.nlm.nih.gov/27160594/)
 24. K. Halldórsdóttir, E. Árnason, Trans-species polymorphism at antimicrobial innate immunity cathelicidin genes of Atlantic cod and related species. *PeerJ* **3**, e976 (2015). doi: [10.7717/peerj.976](https://doi.org/10.7717/peerj.976); pmid: [26038731](https://pubmed.ncbi.nlm.nih.gov/26038731/)
 25. O. Heldgren, B. C. Sheldon, Locus-specific protocol for nine different innate immune genes (antimicrobial peptides: B-defensins) across passerine bird species reveals within-species coding variation and a case of trans-species polymorphisms. *Mol. Ecol. Resour.* **11**, 686–692 (2011). doi: [10.1111/j.1755-0998.2011.02995.x](https://doi.org/10.1111/j.1755-0998.2011.02995.x); pmid: [21676198](https://pubmed.ncbi.nlm.nih.gov/21676198/)
 26. R. M. Linzmeier, T. Ganz, Human defensin gene copy number polymorphisms: Comprehensive analysis of independent variation in α - and β -defensin regions at 8p22-p23. *Genomics* **86**, 423–430 (2005). doi: [10.1016/j.ygeno.2005.06.003](https://doi.org/10.1016/j.ygeno.2005.06.003); pmid: [16039093](https://pubmed.ncbi.nlm.nih.gov/16039093/)
 27. K. J. Smith, E. Gwyer Findlay, Expression of antimicrobial host defence peptides in the central nervous system during health and disease. *Discov. Immunol.* **1**, kyac003 (2022). doi: [10.1093/discim/kyac003](https://doi.org/10.1093/discim/kyac003)
 28. M. A. Hanson, B. Lemaître, Repeated truncation of a modular antimicrobial peptide gene for neural control. *PLoS Genet.* **18**, e1010259 (2022). doi: [10.1371/journal.pgen.1010259](https://doi.org/10.1371/journal.pgen.1010259); pmid: [35714143](https://pubmed.ncbi.nlm.nih.gov/35714143/)
 29. J.-P. Parvy *et al.*, The antimicrobial peptide defensin cooperates with tumour necrosis factor to drive tumour cell death in *Drosophila*. *eLife* **8**, e45061 (2019). doi: [10.7554/eLife.45061](https://doi.org/10.7554/eLife.45061); pmid: [31358113](https://pubmed.ncbi.nlm.nih.gov/31358113/)
 30. S. Lee *et al.*, Downregulation of Hsp90 and the antimicrobial peptide Mtk suppresses poly(GR)-induced neurotoxicity in C9ORF72-ALS/FTD. *Neuron* **111**, 1381–1390.e6 (2023). doi: [10.1016/j.neuron.2023.02.029](https://doi.org/10.1016/j.neuron.2023.02.029); pmid: [36931278](https://pubmed.ncbi.nlm.nih.gov/36931278/)
 31. R. Barajas-Azpeleta *et al.*, Antimicrobial peptides modulate long-term memory. *PLoS Genet.* **14**, e1007440 (2018). pmid: [30312294](https://pubmed.ncbi.nlm.nih.gov/30312294/)
 32. I. S. Pais, R. S. Valente, M. Sporniak, L. Teixeira, *Drosophila melanogaster* establishes a species-specific mutualistic interaction with stable gut-colonizing bacteria. *PLoS Biol.* **16**, e2005710 (2018). doi: [10.1371/journal.pbio.2005710](https://doi.org/10.1371/journal.pbio.2005710); pmid: [29975680](https://pubmed.ncbi.nlm.nih.gov/29975680/)
 33. D. R. Sannino, A. J. Dobson, K. Edwards, E. R. Angert, N. Buchon, The *Drosophila melanogaster* gut microbiota provisions thiamine to its host. *mBio* **9**, e00155–e18 (2018). doi: [10.1128/mBio.00155-18](https://doi.org/10.1128/mBio.00155-18); pmid: [29511074](https://pubmed.ncbi.nlm.nih.gov/29511074/)
 34. J.-S. Chen, S.-C. Tsaur, C.-T. Ting, S. Fang, Dietary utilization drives the differentiation of gut bacterial communities between specialist and generalist drosophilid flies. *Microbiol. Spectr.* **10**, e0141822 (2022). doi: [10.1128/spectrum.01418-22](https://doi.org/10.1128/spectrum.01418-22); pmid: [35863034](https://pubmed.ncbi.nlm.nih.gov/35863034/)
 35. J. Consuegra *et al.*, *Drosophila*-associated bacteria differentially shape the nutritional requirements of their host during juvenile growth. *PLoS Biol.* **18**, e3000681 (2020). doi: [10.1371/journal.pbio.3000681](https://doi.org/10.1371/journal.pbio.3000681); pmid: [32196485](https://pubmed.ncbi.nlm.nih.gov/32196485/)
 36. A. Marra, M. A. Hanson, S. Kondo, B. Erkosar, B. Lemaître, *Drosophila* antimicrobial peptides and lysozymes regulate gut microbiota composition and abundance. *mBio* **12**, e0082421 (2021). doi: [10.1101/2021.03.19.436153](https://doi.org/10.1101/2021.03.19.436153); pmid: [34253067](https://pubmed.ncbi.nlm.nih.gov/34253067/)
 37. S. J. Perlman, J. Jaenike, Infection success in novel hosts: An experimental and phylogenetic study of *Drosophila*-parasitic nematodes. *Evolution* **57**, 544–557 (2003). doi: [10.1111/j.0014-3820.2003.tb01546.x](https://doi.org/10.1111/j.0014-3820.2003.tb01546.x); pmid: [12703944](https://pubmed.ncbi.nlm.nih.gov/12703944/)
 38. Y. Carton, M. Bouletreau, J. van Alphen, J. C. van Lenteren, “The *Drosophila* parasitic wasps,” in *The Genetics and Biology of Drosophila* (Academic, 1986), vol. 3, pp. 348–394.
 39. A. N. Brown, V. K. Lloyd, Evidence for horizontal transfer of Wolbachia by a *Drosophila* mite. *Exp. Appl. Acarol.* **66**, 301–311 (2015). doi: [10.1007/s10493-015-9918-z](https://doi.org/10.1007/s10493-015-9918-z); pmid: [25921489](https://pubmed.ncbi.nlm.nih.gov/25921489/)
 40. I. Kounatidis *et al.*, *Acetobacter tropicalis* is a major symbiont of the olive fruit fly (*Bactrocera oleae*). *Appl. Environ. Microbiol.* **75**, 3281–3288 (2009). doi: [10.1128/AEM.02933-08](https://doi.org/10.1128/AEM.02933-08); pmid: [19304418](https://pubmed.ncbi.nlm.nih.gov/19304418/)
 41. J. A. Chandler, J. M. Lang, S. Bhatnagar, J. A. Eisen, A. Kopp, Bacterial communities of diverse *Drosophila* species: Ecological context of a host-microbe model system. *PLoS Genet.* **7**, e1002272 (2011). doi: [10.1371/journal.pgen.1002272](https://doi.org/10.1371/journal.pgen.1002272); pmid: [21966276](https://pubmed.ncbi.nlm.nih.gov/21966276/)
 42. J. L. Dimarcq *et al.*, Insect immunity. Purification and characterization of a family of novel inducible antibacterial proteins from immunized larvae of the dipteran *Phormia terranova* and complete amino-acid sequence of the predominant member, dipterocin A. *Eur. J. Biochem.* **171**, 17–22 (1988). doi: [10.1111/j.1432-1033.1988.tb13752.x](https://doi.org/10.1111/j.1432-1033.1988.tb13752.x); pmid: [3276515](https://pubmed.ncbi.nlm.nih.gov/3276515)
 43. M. A. Hanson, B. Lemaître, R. L. Unckless, Dynamic evolution of antimicrobial peptides underscores trade-offs between immunity and ecological fitness. *Front. Immunol.* **10**, 2620 (2019). doi: [10.3389/fimmu.2019.02620](https://doi.org/10.3389/fimmu.2019.02620); pmid: [31781114](https://pubmed.ncbi.nlm.nih.gov/31781114)
 44. M. A. Hanson, P. T. Hamilton, S. J. Perlman, Immune genes and divergent antimicrobial peptides in flies of the subgenus *Drosophila*. *BMC Evol. Biol.* **16**, 228 (2016). doi: [10.1186/s12862-016-0805-y](https://doi.org/10.1186/s12862-016-0805-y); pmid: [27776480](https://pubmed.ncbi.nlm.nih.gov/27776480)
 45. A. Suvorov *et al.*, Widespread introgression across a phylogeny of 155 *Drosophila* genomes. *Curr. Biol.* **32**, 111–123.e5 (2021). doi: [10.1016/j.cub.2021.10.052](https://doi.org/10.1016/j.cub.2021.10.052); pmid: [34788634](https://pubmed.ncbi.nlm.nih.gov/34788634)
 46. X. Bing, J. Gerlach, G. Loeb, N. Buchon, Nutrient-dependent impact of microbes on *Drosophila suzukii* development. *mBio* **9**, e02199–e17 (2018). doi: [10.1128/mBio.02199-17](https://doi.org/10.1128/mBio.02199-17); pmid: [29559576](https://pubmed.ncbi.nlm.nih.gov/29559576)
 47. I. Martínez-Sañudo *et al.*, Metagenomic analysis reveals changes of the *Drosophila suzukii* microbiota in the newly colonized regions. *Insect Sci.* **25**, 833–846 (2018). doi: [10.1111/1744-7917.12458](https://doi.org/10.1111/1744-7917.12458); pmid: [28323391](https://pubmed.ncbi.nlm.nih.gov/28323391)
 48. V. G. Martinson, A. E. Douglas, J. Jaenike, Community structure of the gut microbiota in sympatric species of wild *Drosophila*. *Ecol. Lett.* **20**, 629–639 (2017). doi: [10.1111/ele.12761](https://doi.org/10.1111/ele.12761); pmid: [28371064](https://pubmed.ncbi.nlm.nih.gov/28371064)
 49. A. Bost *et al.*, Functional variation in the gut microbiome of wild *Drosophila* populations. *Mol. Ecol.* **27**, 2834–2845 (2018). doi: [10.1111/mec.14728](https://doi.org/10.1111/mec.14728); pmid: [29802796](https://pubmed.ncbi.nlm.nih.gov/29802796)
 50. B. Goldman-Huertas *et al.*, Evolution of herbivory in Drosophilidae linked to loss of behaviors, antennal receptors, odorant receptors, and ancestral diet. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 3026–3031 (2015). doi: [10.1073/pnas.1424656112](https://doi.org/10.1073/pnas.1424656112); pmid: [25624509](https://pubmed.ncbi.nlm.nih.gov/25624509)
 51. T. K. O'Connor, P. T. Humphrey, R. T. Lapoint, N. K. Whiteman, P. M. O'Grady, Microbial interactions and the ecology and evolution of Hawaiian Drosophilidae. *Front. Microbiol.* **5**, 616 (2014). pmid: [25566196](https://pubmed.ncbi.nlm.nih.gov/25566196)
 52. R. R. Copley *et al.*, Functional conservation of Rel binding sites in drosophilid genomes. *Genome Res.* **17**, 1327–1335 (2007). doi: [10.1101/gr.6490707](https://doi.org/10.1101/gr.6490707); pmid: [17785540](https://pubmed.ncbi.nlm.nih.gov/17785540)
 53. A. D. Gloss *et al.*, Evolution of herbivory remodels a *Drosophila* genome. *bioRxiv* 767160 [Preprint] (2019); <https://doi.org/10.1101/767160>
 54. M. De Cock *et al.*, Comparative microbiomics of tephritid frugivorous pests (Diptera: Tephritidae) from the field: A tale of high variability across and within species. *Front. Microbiol.* **11**, 1890 (2020). doi: [10.3389/fmicb.2020.01890](https://doi.org/10.3389/fmicb.2020.01890); pmid: [32849469](https://pubmed.ncbi.nlm.nih.gov/32849469)
 55. L. Mazzon *et al.*, Presence of specific symbiotic bacteria in flies of the subfamily Tephritinae (Diptera Tephritidae) and their phylogenetic relationships: Proposal of ‘Candidatus Stammerula tephritidis’. *Int. J. Syst. Evol. Microbiol.* **58**, 1277–1287 (2008). doi: [10.1099/ijs.0.65287-0](https://doi.org/10.1099/ijs.0.65287-0); pmid: [18523165](https://pubmed.ncbi.nlm.nih.gov/18523165)
 56. N. Mollentze, D. Keen, U. Munkhbayar, R. Biek, D. G. Streicker, Variation in the ACE2 receptor has limited utility for SARS-CoV-2 host prediction. *eLife* **11**, e80329 (2022). doi: [10.7554/eLife.80329](https://doi.org/10.7554/eLife.80329); pmid: [36416537](https://pubmed.ncbi.nlm.nih.gov/36416537)
 57. R. M. Imrie, K. E. Roberts, B. Longdon, Between virus correlations in the outcome of infection across host species: Evidence of virus genotype by host species interactions. *Evol. Lett.* **5**, 472–483 (2021). doi: [10.1101/2021.02.16.431403](https://doi.org/10.1101/2021.02.16.431403); pmid: [34621534](https://pubmed.ncbi.nlm.nih.gov/34621534)
 58. M. Moraitou *et al.*, Ecology, not host phylogeny, shapes the oral microbiome in closely related species. *Mol. Biol. Evol.* **39**, msac263 (2022). doi: [10.1093/molbev/msac263](https://doi.org/10.1093/molbev/msac263); pmid: [36472532](https://pubmed.ncbi.nlm.nih.gov/36472532)
 59. F. H. Login *et al.*, Antimicrobial peptides keep insect endosymbionts under control. *Science* **334**, 362–365 (2011). doi: [10.1126/science.1209728](https://doi.org/10.1126/science.1209728); pmid: [22021855](https://pubmed.ncbi.nlm.nih.gov/22021855)
 60. N. M. Gerardo *et al.*, Immunity and other defenses in pea aphids, *Acyrthosiphon pisum*. *Genome Biol.* **11**, R21 (2010). doi: [10.1186/gb-2010-11-2-r21](https://doi.org/10.1186/gb-2010-11-2-r21); pmid: [20178569](https://pubmed.ncbi.nlm.nih.gov/20178569)
 61. C. Vörburger, S. J. Perlman, The role of defensive symbionts in host-parasite coevolution. *Biol. Rev. Camb. Philos. Soc.* **93**, 1747–1764 (2018). doi: [10.1111/brv.12417](https://doi.org/10.1111/brv.12417); pmid: [29663622](https://pubmed.ncbi.nlm.nih.gov/29663622)
 62. P. Bulet *et al.*, Insect immunity. The inducible antibacterial peptide dipterocin carries two O-glycans necessary for biological activity. *Biochemistry* **34**, 7394–7400 (1995). doi: [10.1021/bi00022a012](https://doi.org/10.1021/bi00022a012); pmid: [7779781](https://pubmed.ncbi.nlm.nih.gov/7779781)
 63. M. Cudic, P. Bulet, R. Hoffmann, D. J. Craik, L. Otvos Jr., Chemical synthesis, antibacterial activity and conformation of dipterocin, an 82-mer peptide originally isolated from insects. *Eur. J. Biochem.* **266**, 549–558 (1999). doi: [10.1046/j.1432-1327.1999.00894.x](https://doi.org/10.1046/j.1432-1327.1999.00894.x); pmid: [10561597](https://pubmed.ncbi.nlm.nih.gov/10561597)
 64. A. D. Long, S. J. Macdonald, E. G. King, Dissecting complex traits using the *Drosophila* Synthetic Population Resource. *Trends Genet.* **30**, 488–495 (2014). doi: [10.1016/j.tig.2014.07.009](https://doi.org/10.1016/j.tig.2014.07.009); pmid: [25175100](https://pubmed.ncbi.nlm.nih.gov/25175100)
 65. B. Vicoso, D. Bachtrog, Numerous transitions of sex chromosomes in Diptera. *PLoS Biol.* **13**, e1002078 (2015). doi: [10.1371/journal.pbio.1002078](https://doi.org/10.1371/journal.pbio.1002078); pmid: [25879221](https://pubmed.ncbi.nlm.nih.gov/25879221)
 66. C. H. Scott Chialvo, B. E. White, L. K. Reed, K. A. Dyer, A phylogenetic examination of host use evolution in the quinaria and testacea groups of *Drosophila*. *Mol. Phylogenet. Evol.* **130**, 233–243 (2019). doi: [10.1016/j.ympev.2018.10.027](https://doi.org/10.1016/j.ympev.2018.10.027); pmid: [30366088](https://pubmed.ncbi.nlm.nih.gov/30366088)
 67. M. A. Khallaf *et al.*, Large-scale characterization of sex pheromone communication systems in *Drosophila*. *Nat. Commun.* **12**, 4165 (2021). doi: [10.1101/2020.09.21.305854](https://doi.org/10.1101/2020.09.21.305854); pmid: [34230464](https://pubmed.ncbi.nlm.nih.gov/34230464)
 68. Data for: M. A. Hanson, L. Grollmus, B. Lemaître, Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*. *Dryad* (2023); doi: [10.5061/dryad.dz08kps2p](https://doi.org/10.5061/dryad.dz08kps2p)

ACKNOWLEDGMENTS

We thank members of N. Whiteman's laboratory, as well as V. Martinson and B. Longdon for consultation; S. Macdonald for the *DptB*^{A3} stock; J. P. Boquete for transgenic fly injections; and other colleagues for comments on previous versions of this manuscript.

Funding: This work was funded by the Swiss National Science Foundation (SNSF postdoctoral mobility grant P500PB_211082 to M.A.H.), Sinergia (grant CRSII5_186397 to B.L.), and the Novartis Foundation (grant 532114 to B.L.). **Author contributions:** M.A.H. and B.L. designed the study. M.A.H. conceived experiments and designed fly genetics. M.A.H. and L.G. performed experiments. L.G. provided essential molecular and experimental support. M.A.H. performed bioinformatic and literature survey analyses. B.L. supervised this work. M.A.H. and B.L. wrote the manuscript.

Competing interests: The authors declare no competing interests. **Data and materials availability:** All data generated in this study, including tables with all experimental raw data, A. sicerae BELCH sequencing statistics, and alignments and phylogenies, are given in nested folder format sorted per relevant figure and table at Dryad (68). Certain fly and bacteria stocks came from the Bloomington *Drosophila* Resource Centre (<https://bdsc.indiana.edu/information/patent-license.html>) or the Leibniz Institute DSMZ collection of microorganisms (<https://www.dsmz.de/collection/nagoya-protocol/legal-background>) and are described in the supplementary materials and methods. All other fly and bacteria stocks are freely available upon request to B.L. We thank Uniprot (<https://www.uniprot.org/>) and FlyBase (<https://flybase.org/>) for database management. **License information:** Copyright © 2023 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.science.org/about/science-licenses-journal-article-reuse>

SUPPLEMENTARY MATERIALS

science.org/doi/10.1126/science.adg5725

Materials and Methods

Figs. S1 to S17

Tables S1 to S4

References (69–81)

MDAR Reproducibility Checklist

Submitted 6 January 2023; accepted 8 June 2023

10.1126/science.adg5725



Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*

M. A. Hanson, L. Grollmus, and B. Lemaitre

Science, **381** (6655), eadg5725.

DOI: 10.1126/science.adg5725

Editor's summary

To prevent gut microbiota from running amok, animals and plants secrete a series of small, often multifunctional peptides called antimicrobial peptides. Until recently, antimicrobial peptides were considered to have broad activities, and it was unclear why such molecules showed signs of rapid evolution. Hanson *et al.* found a striking specificity for the peptides diptericin A and B for two species of gut commensal bacteria. These species occur in the natural environment of fruit flies depending on the food resource exploited: fruit or fungi. Thus, the presence or absence of diptericin A or B predicts the ecology of the fly. This work shows how an organism's microbiota might be able to shape the host's immune responses in a manner similar to how a host's immune responses shape its microbiota. —Caroline Ash

View the article online

<https://www.science.org/doi/10.1126/science.adg5725>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

Science (ISSN) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works